

ENGINEERING  
LIBRARY

CIT - ELECTRON TUBE & MICROWAVE  
LABORATORY REPORT

ECHOES IN COLLISION-FREE PLASMAS

R. W. Gould

Technical Report No. 36

July 1967

CALIFORNIA INSTITUTE OF TECHNOLOGY

PASADENA, CALIFORNIA

CALIFORNIA  
INSTITUTE OF

OCT 16 1975

TECHNOLOGY

ECHOES IN COLLISION-FREE PLASMAS

R. W. Gould

Technical Report No. 36

July 1967

A report on research sponsored by the

Office of Naval Research

Contract Nonr 220(50)

(Submitted to Physics Letters for publication)

California Institute of Technology

Pasadena, California

# ECHOES IN COLLISION-FREE PLASMAS

R. W. Gould

California Institute of Technology, Pasadena, California

## ABSTRACT

It is shown that temporal echoes can occur in collision-free plasmas at various times for different wave number combinations and that spatial echoes can occur at various positions for different frequency combinations.

A plasma wave echo in collision-free plasmas has been predicted from a perturbation solution of the Vlasov equation<sup>1</sup>. Here we demonstrate the existence of higher order temporal as well as spatial echoes, based on a simple nonperturbation method which gives results to all orders. The self-consistent field of the plasma is, however, neglected.

We consider a plasma of charged particles subjected to two external electric field pulses with different wave numbers at  $t = 0, \tau$ , i.e.  $qE_x/M = v_1 \delta(t) \cos k_1 x + v_2 \delta(t-\tau) \cos k_2 x$ . The position of a charged particle whose initial position and velocity is  $(x_0, u_0)$  in such a field is readily found to be

$$\begin{aligned} x(t, x_0, u_0) &= x_0 + t(u_0 + v_1 \cos[k_1 x_0]) & 0 \leq t \leq \tau \\ &= x(\tau, x_0, u_0) + v_2(t-\tau) \cos[k_2 x(\tau, x_0, u_0)] & \tau \leq t < \infty \end{aligned} \quad (1)$$

The spatial Fourier components of the charge density may be obtained by integrating over the initial (Maxwellian) distribution  $F_0(x_0, u_0)$ ,  $\rho_k(t) = q \iint F_0(x_0, u_0) \exp[-ikx(t, x_0, u_0)] dx_0 du_0$ . Expanding the exponential using well known Bessel function identities<sup>2</sup> and performing the integrations, we find that  $\rho_k(t)$  vanishes unless

$$k = -mk_1 + nk_2 \equiv k_{mn} \quad (2)$$

whereupon

$$\rho_k(t) = n_0 q A_{mn}(t) g_{mn}(t - \tau_{mn}) \quad (3)$$

with

$$\tau_{mn} = (nk_2/k_{mn}) \tau \quad (4)$$

$$g_{mn}(t) = (1/m!)(k_{mn} \bar{v} t/2)^m e^{-(k_{mn} \bar{v} t/2)^2} \quad (5)$$

$$A_{mn}(t) = (-i)^{n-m} \frac{J_m[k_{mn} v_1(t-\tau_{mn})]}{[k_{mn} \bar{v}(t-\tau_{mn})/2]^m/m!} J_n[k_{mn} v_2(t-\tau)] \quad (6)$$

$n_0$  is the average density of charged particles and  $\bar{v} = \sqrt{2kT/M}$  is their r.m.s. thermal speed. When  $v_1$  and  $v_2$  are small compared with  $\bar{v}$ ,  $A_{mn}(t)$  is slowly varying compared to  $g_{mn}(t)$ . Furthermore, only  $A_{mn}(t)$  depends on the pulse amplitudes  $v_1$  and  $v_2$ . In this limit  $A_{mn}(\tau_{mn})$  gives the amplitude of the pulse and  $g_{mn}(t)$  gives its shape. Thus we see that, provided  $\tau_{mn} > \tau$ , each spatial Fourier component with combination wave number given by (2) exhibits a temporal echo at time  $\tau_{mn}$  which is determined by the separation of the two applied pulses and the ratio of their wave numbers. For weak pulses ( $v_1$  and  $v_2$  small) the amplitude reduces to

$$A_{mn} = (-i)^{n-m} \left(\frac{v_1}{\bar{v}}\right)^m \left(\frac{v_2}{\bar{v}}\right)^n \frac{1}{n!} \left(\frac{mk_1 \bar{v} \tau}{2}\right)^n \quad (7)$$

which exhibits the expected power law dependence on the pulse amplitudes  $v_1$  and  $v_2$  and the pulse separation  $\tau$ .

A similar analysis has been made for spatial echoes, which may be more important experimentally. When spatially localized electric fields of differing frequencies  $\omega_1$  and  $\omega_2$  are applied at  $z = 0$  and  $z = l$  respectively, in the plasma,  $E_x = V_1 \delta(x) \cos \omega_1 t + V_2 \delta(x-l) \cos \omega_2 t$ . The temporal Fourier components of the electric current density are found



to vanish unless

$$\omega = -m\omega_1 + n\omega_2 \equiv \omega_{mn} \quad (8)$$

and that each of these exhibits a localized maximum at the position

$$x = (n\omega_2/\omega_{mn})\ell \equiv \ell_{mn} \quad (9)$$

provided that  $\ell_{mn} > \ell$ . We also find for the amplitude

$$A_{mn}(\ell_{mn}) \sim \left[ \frac{m\omega_1 \ell}{2v} \frac{eV_2}{Mv^2} \right]^n \left[ \frac{n\omega_2 \ell}{2v} \frac{eV_1}{Mv^2} \right]^m \quad (10)$$

when the dimensionless quantities in the brackets are small.

(8), (9), and (10) are the spatial analogs of (2), (4) and (6), respectively. Thus each temporal Fourier component of the electric current with frequencies given by (8) exhibits a spatial echo at a position which is determined by (9).

In obtaining these results the self-consistent fields of the plasma have been neglected. This approximation should be valid as long as all wave numbers for which  $\rho_k/\rho_0$  is important are large (all frequencies for which  $i_k/i_0$  is important are large). When these conditions are not satisfied we expect a substantial modification in the echo amplitudes and shapes. However, we still expect (4) [or (9)] to correctly give the time [or position] of the echoes of the appropriate wave number (2) [or frequency (8)]. In addition, for small amplitudes we still expect the power law dependence on  $v_1, v_2$  and  $\tau$  [or  $V_1, V_2$  and  $\ell$ ] suggested by (6) [or (10)] to hold.

The author is indebted to T. M. O'Neil and J. H. Malmberg for helpful discussions and to the Office of Naval Research for its sponsorship of this research.

#### References

1. R. W. Gould, T. M. O'Neil, J. H. Malmberg, "The Plasma Wave Echo", Phys. Rev. Letters (to appear).
2. W. Magnus, F. Oberhettinger, Special Functions of Mathematical Physics, Chelsea 1949, New York, see p. 18 and p. 20.

UNCLASSIFIED

Security Classification

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

## 1. ORIGINATING ACTIVITY (Corporate author)

California Institute of Technology  
Pasadena, California 91109

## 2a. REPORT SECURITY CLASSIFICATION

UNCLASSIFIED

## 2b. GROUP

## 3. REPORT TITLE

ECHOES IN COLLISION-FREE PLASMAS

## 4. DESCRIPTIVE NOTES (Type of report and inclusive dates)

Scientific Interim

## 5. AUTHOR(S) (First name, middle initial, last name)

Roy W. Gould

## 6. REPORT DATE

July 1967

## 7a. TOTAL NO. OF PAGES

6

## 7b. NO. OF REFS

2

## 8a. CONTRACT OR GRANT NO.

Nonr 220(50)

## b. PROJECT NO.

c.

d.

## 9a. ORIGINATOR'S REPORT NUMBER(S)

Tech. Report No. 36

## 9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)

## 10. DISTRIBUTION STATEMENT

DISTRIBUTION OF THIS DOCUMENT IS UNLIMITED

## 11. SUPPLEMENTARY NOTES

## 12. SPONSORING MILITARY ACTIVITY

Electronics Division 427  
Office of Naval Research  
Washington, D. C.

## 13. ABSTRACT

It is shown that temporal echoes can occur in collision-free plasmas at various times for different wave number combinations and that spatial echoes can occur at various positions for different frequency combinations.

DD FORM 1473  
1 NOV 65

UNCLASSIFIED

Security Classification



UNCLASSIFIED